

**NICAPRIN™**  
**Rx Only Dietary Supplement**

**DESCRIPTION:**

This product is a prescription dietary supplement for oral administration, specifically formulated for the dietary management of patients with unique nutritional needs who require increased levels of one or more of the ingredients in this product. Each tan colored caplet-shaped tablet is debossed with "203" on one side and plain on the other.

**Supplement Facts**

Serving Size: 1 Tablet

Amount Per Serving	
Niacin (nicotinamide)	250 mg
Folic Acid	500 mcg
Vitamin B <sub>6</sub> (pyridoxine HCl)	4.5 mg
Copper (cupric oxide)	1.4 mg
Zinc (zinc oxide)	9 mg
Soliprin (scutellaria baicalensis (root) & acacia catechu (heartwood))	125 mg

**OTHER INGREDIENTS:** Iron oxide, magnesium stearate, methacrylic acid copolymer, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, sodium bicarbonate, starch, silica, talc, and titanium dioxide.

**INDICATIONS AND USAGE:** This product is indicated for patients who are deficient in, or at risk of deficiency in, one or more of the components of this product.

**CONTRAINDICATIONS:** This product is contraindicated in patients with known hypersensitivity to any of the ingredients.

**WARNINGS AND PRECAUTIONS:** This product should be administered with caution in patients with a history of liver disease, jaundice or diabetes mellitus. While prescribing this product for pregnant women, nursing mothers, or for women prior to conception, their medical condition and

use of other drugs, herbs and/or supplements should be considered.

Folic acid above 1 mg daily may obscure pernicious anemia in that hematologic remission may occur while neurological manifestations remain progressive. Abnormal liver functions tests have been reported in persons taking daily doses of 500 mg or more of niacinamide.

**KEEP THIS AND ALL MEDICATION OUT OF THE REACH OF CHILDREN.**

**PREGNANCY & NURSING MOTHERS:** This product is not indicated for use as a prenatal/postnatal multivitamin for lactating and nonlactating mothers. Physicians and medical practitioners should administer this product with caution to patients who are pregnant, lactating and/or taking medication.

**ADVERSE REACTIONS:** Allergic sensitization has been reported following both oral and parenteral administration of folic acid. **You should call your doctor for medical advice about side effects. To report a serious adverse event, call 1-855-899-4237.**

**DOSAGE AND ADMINISTRATION:** Usual adult dose is 1 to 2 tablets daily, or as directed by a physician.

**HOW SUPPLIED:** This product is supplied in the following size(s):  
60 count bottles, **NDC 57893-203-60**

Store between 15°C and 30°C (between 59°F and 86°F). Tamper evident by foil seal under cap.

Manufactured for:  
Artesa Labs, LLC  
137585 Research Blvd.  
Suite 125  
Austin, TX 78750

v1 Rev. 05/16 826146

Artesa Labs